

Ce site utilise des cookies provenant de Google pour fournir ses services et analyser le trafic. Votre adresse IP et votre user-agent, ainsi que des statistiques relatives aux performances et à la sécurité, sont transmis à Google afin d'assurer un service de qualité, de générer des statistiques d'utilisation, et de détecter et de résoudre les problèmes d'abus.

EN SAVOIR PLUS **OK !**

```

00 0000: 0 (Atom)

```

Share this

About Me

Total Pageviews

Share This

Powered by Blogger.

[REDACTED]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to the quality of the scan and the density of the text. The text appears to be a single paragraph or a series of closely related sentences. The visible fragments of text are as follows:]

[Illegible text block]

[REDACTED]

[REDACTED]

the 1990s, the number of people in the UK who are aged 65 and over has increased by 1.5 million, and the number of people aged 75 and over has increased by 1.2 million (Office for National Statistics 2000). The number of people aged 85 and over has increased by 0.5 million in the same period.

There is a growing awareness of the need to develop services to meet the needs of the ageing population. The Department of Health (1999) has set out a strategy for the future of health care for older people. The strategy is based on the principle that older people should be able to live in their own homes for as long as possible, and that health care should be provided in a way that is appropriate to their needs. The strategy is based on the following principles:

- Older people should be able to live in their own homes for as long as possible.
- Health care should be provided in a way that is appropriate to the needs of older people.
- Older people should be able to access the services they need when they need them.
- Older people should be able to participate in decisions about their care.

The strategy is based on the principle that older people should be able to live in their own homes for as long as possible, and that health care should be provided in a way that is appropriate to their needs. The strategy is based on the following principles:

- Older people should be able to live in their own homes for as long as possible.
- Health care should be provided in a way that is appropriate to the needs of older people.
- Older people should be able to access the services they need when they need them.
- Older people should be able to participate in decisions about their care.

The strategy is based on the principle that older people should be able to live in their own homes for as long as possible, and that health care should be provided in a way that is appropriate to their needs. The strategy is based on the following principles:

- Older people should be able to live in their own homes for as long as possible.
- Health care should be provided in a way that is appropriate to the needs of older people.
- Older people should be able to access the services they need when they need them.
- Older people should be able to participate in decisions about their care.

The strategy is based on the principle that older people should be able to live in their own homes for as long as possible, and that health care should be provided in a way that is appropriate to their needs. The strategy is based on the following principles:

- Older people should be able to live in their own homes for as long as possible.
- Health care should be provided in a way that is appropriate to the needs of older people.
- Older people should be able to access the services they need when they need them.
- Older people should be able to participate in decisions about their care.

The strategy is based on the principle that older people should be able to live in their own homes for as long as possible, and that health care should be provided in a way that is appropriate to their needs. The strategy is based on the following principles:

- Older people should be able to live in their own homes for as long as possible.
- Health care should be provided in a way that is appropriate to the needs of older people.
- Older people should be able to access the services they need when they need them.
- Older people should be able to participate in decisions about their care.

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[REDACTED]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of closely related sentences, but the specific words and structure cannot be discerned.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of closely related sentences, but the specific words and structure cannot be discerned.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It appears to be a mix of English and possibly some non-English characters, but the overall structure suggests a single paragraph or a series of closely related sentences. The text is too blurry and low-contrast to transcribe accurately, but it seems to contain several lines of prose.]

[Illegible text block]

[The following text is a dense, continuous block of text, likely a scan of a document page. It appears to be a mix of English and possibly some non-English characters, but the overall structure suggests a single paragraph or a series of lines of text. Due to the low resolution and potential noise in the scan, the specific words and punctuation are difficult to discern accurately. The text is oriented vertically on the page.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to the quality of the scan and the density of the text. The text appears to be a single paragraph or a series of closely related sentences. The visible fragments of text are as follows:]

[Illegible text block]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of closely related sentences, but the specific words and structure cannot be discerned.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It appears to be a mix of English and possibly some non-English characters, but the overall structure suggests a single paragraph or a series of lines of text. Due to the low resolution and potential noise in the scan, the specific words and punctuation are difficult to discern accurately. The text is oriented vertically on the page.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of closely related sentences, but the specific words and structure cannot be discerned.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It appears to be a mix of English and possibly some non-English characters, but the overall structure suggests a formal document or report. The text is too blurry and low-contrast to transcribe accurately. It seems to contain several paragraphs of text, possibly including a title or header at the top and a footer at the bottom. The content is illegible due to the quality of the scan.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of closely related sentences, but the specific words and structure cannot be discerned.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to the quality of the scan and the density of the text. The text appears to be a single paragraph or a series of closely related sentences. The visible fragments of text are as follows:]

[Illegible text block]

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported as the most common serotype in children with acute bacterial dysentery [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1980s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [12]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [13].

The purpose of this study was to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low resolution. The text appears to be a single paragraph or a section of a larger document, but the specific words and sentences cannot be transcribed accurately.]

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported as the most common serotype in children with acute bacterial dysentery [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. The only published study of *S. flexneri* in the United Kingdom was by Smith *et al.* [12], who reported that *S. flexneri* was the most common serotype isolated from children with acute bacterial dysentery in the United Kingdom in 1987. The serotypes isolated were *S. flexneri* 3, 4, 5, 6, 7, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100.

The aim of this study was to determine the prevalence of *S. flexneri* in children with acute bacterial dysentery in the United Kingdom in 1997. The study was part of a larger study of the epidemiology of acute bacterial dysentery in the United Kingdom in 1997, which was conducted by the Health Protection Agency (HPA) and the Public Health Laboratory Service (PHLS).

The study was conducted in 1997 in the United Kingdom. The study was part of a larger study of the epidemiology of acute bacterial dysentery in the United Kingdom in 1997, which was conducted by the Health Protection Agency (HPA) and the Public Health Laboratory Service (PHLS).

The study was conducted in 1997 in the United Kingdom. The study was part of a larger study of the epidemiology of acute bacterial dysentery in the United Kingdom in 1997, which was conducted by the Health Protection Agency (HPA) and the Public Health Laboratory Service (PHLS).

The study was conducted in 1997 in the United Kingdom. The study was part of a larger study of the epidemiology of acute bacterial dysentery in the United Kingdom in 1997, which was conducted by the Health Protection Agency (HPA) and the Public Health Laboratory Service (PHLS).

The study was conducted in 1997 in the United Kingdom. The study was part of a larger study of the epidemiology of acute bacterial dysentery in the United Kingdom in 1997, which was conducted by the Health Protection Agency (HPA) and the Public Health Laboratory Service (PHLS).

The study was conducted in 1997 in the United Kingdom. The study was part of a larger study of the epidemiology of acute bacterial dysentery in the United Kingdom in 1997, which was conducted by the Health Protection Agency (HPA) and the Public Health Laboratory Service (PHLS).

The study was conducted in 1997 in the United Kingdom. The study was part of a larger study of the epidemiology of acute bacterial dysentery in the United Kingdom in 1997, which was conducted by the Health Protection Agency (HPA) and the Public Health Laboratory Service (PHLS).

The study was conducted in 1997 in the United Kingdom. The study was part of a larger study of the epidemiology of acute bacterial dysentery in the United Kingdom in 1997, which was conducted by the Health Protection Agency (HPA) and the Public Health Laboratory Service (PHLS).

The study was conducted in 1997 in the United Kingdom. The study was part of a larger study of the epidemiology of acute bacterial dysentery in the United Kingdom in 1997, which was conducted by the Health Protection Agency (HPA) and the Public Health Laboratory Service (PHLS).

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported as the most common serotype in children with acute bacterial dysentery [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1980s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [12]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [13]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [15].

In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [16]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [17]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [18]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [19].

In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [20]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [21]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [22]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [23].

In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [24]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [25]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [26]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [27].

In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [28]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [29]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [30]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [31].

In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [32]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [33]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [34]. In the 1990s, *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom [35].

