

Ce site utilise des cookies provenant de Google pour fournir ses services et analyser le trafic. Votre adresse IP et votre user-agent, ainsi que des statistiques relatives aux performances et à la sécurité, sont transmis à Google afin d'assurer un service de qualité, de générer des statistiques d'utilisation, et de détecter et de résoudre les problèmes d'abus.

EN SAVOIR PLUS **OK !**

```
00 0000: 0 (Atom)
```

Share this

About Me



Total Pageviews

Share This

Powered by Blogger.

[The following text is a dense, continuous block of characters and symbols, likely representing a corrupted or heavily redacted document. It contains no legible words or phrases.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of closely related sentences, but the specific words and structure cannot be discerned.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to the quality of the scan and the density of the text. The text appears to be a single paragraph or a series of closely related sentences. The visible fragments of text are as follows:]

[Illegible text block]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to the quality of the scan and the density of the text. The text appears to be a single paragraph or a series of closely related sentences. The visible fragments of text are as follows:]

[Illegible text block]

[The following text is a dense, continuous block of characters and symbols, likely representing a corrupted or heavily redacted document. It contains no legible words or phrases.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[REDACTED]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to the quality of the scan and the density of the text. The text appears to be a single paragraph or a series of closely related sentences. The visible fragments of text are as follows:]

[Illegible text block]

the 1990s, the number of people in the UK who are employed in the public sector has increased by 1.5 million, from 2.5 million in 1980 to 4 million in 1999. The public sector has also become an important employer of people with disabilities, with 1.5 million people with disabilities employed in the public sector in 1999, compared with 1.2 million in 1980.

There are a number of reasons why the public sector has become an important employer of people with disabilities. One reason is that the public sector has a long history of employing people with disabilities. In the 19th century, the public sector employed people with disabilities in a number of different roles, including as clerks, typists, and stenographers.

Another reason why the public sector has become an important employer of people with disabilities is that the public sector has a number of different departments and agencies, each of which has its own specific needs. This means that the public sector can employ people with disabilities in a wide range of roles, from clerical to professional.

One of the main reasons why the public sector has become an important employer of people with disabilities is that the public sector has a number of different departments and agencies, each of which has its own specific needs. This means that the public sector can employ people with disabilities in a wide range of roles, from clerical to professional.

Another reason why the public sector has become an important employer of people with disabilities is that the public sector has a number of different departments and agencies, each of which has its own specific needs. This means that the public sector can employ people with disabilities in a wide range of roles, from clerical to professional.

One of the main reasons why the public sector has become an important employer of people with disabilities is that the public sector has a number of different departments and agencies, each of which has its own specific needs. This means that the public sector can employ people with disabilities in a wide range of roles, from clerical to professional.

Another reason why the public sector has become an important employer of people with disabilities is that the public sector has a number of different departments and agencies, each of which has its own specific needs. This means that the public sector can employ people with disabilities in a wide range of roles, from clerical to professional.

One of the main reasons why the public sector has become an important employer of people with disabilities is that the public sector has a number of different departments and agencies, each of which has its own specific needs. This means that the public sector can employ people with disabilities in a wide range of roles, from clerical to professional.

Another reason why the public sector has become an important employer of people with disabilities is that the public sector has a number of different departments and agencies, each of which has its own specific needs. This means that the public sector can employ people with disabilities in a wide range of roles, from clerical to professional.

One of the main reasons why the public sector has become an important employer of people with disabilities is that the public sector has a number of different departments and agencies, each of which has its own specific needs. This means that the public sector can employ people with disabilities in a wide range of roles, from clerical to professional.

Another reason why the public sector has become an important employer of people with disabilities is that the public sector has a number of different departments and agencies, each of which has its own specific needs. This means that the public sector can employ people with disabilities in a wide range of roles, from clerical to professional.

One of the main reasons why the public sector has become an important employer of people with disabilities is that the public sector has a number of different departments and agencies, each of which has its own specific needs. This means that the public sector can employ people with disabilities in a wide range of roles, from clerical to professional.

Another reason why the public sector has become an important employer of people with disabilities is that the public sector has a number of different departments and agencies, each of which has its own specific needs. This means that the public sector can employ people with disabilities in a wide range of roles, from clerical to professional.

[REDACTED]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of closely related sentences, but the specific words and structure cannot be discerned.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[REDACTED]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[REDACTED]

the 1990s, the number of people in the UK who are employed in the public sector has increased by 1.5 million, from 2.5 million in 1980 to 4 million in 1999. The public sector has also become an important employer of people with disabilities, with 1.5 million people with disabilities employed in the public sector in 1999, compared with 1.2 million in 1980.

There are a number of reasons why the public sector has become an important employer of people with disabilities. One reason is that the public sector has a long history of employing people with disabilities. In the 19th century, the public sector employed people with disabilities in a number of different roles, including as clerks, typists, and stenographers. In the 20th century, the public sector employed people with disabilities in a number of different roles, including as teachers, nurses, and social workers.

Another reason why the public sector has become an important employer of people with disabilities is that the public sector has a number of policies in place that encourage the employment of people with disabilities. For example, the public sector has a number of policies in place that encourage the employment of people with disabilities in a number of different roles, including as teachers, nurses, and social workers.

There are a number of reasons why the public sector has become an important employer of people with disabilities. One reason is that the public sector has a long history of employing people with disabilities. In the 19th century, the public sector employed people with disabilities in a number of different roles, including as clerks, typists, and stenographers. In the 20th century, the public sector employed people with disabilities in a number of different roles, including as teachers, nurses, and social workers.

Another reason why the public sector has become an important employer of people with disabilities is that the public sector has a number of policies in place that encourage the employment of people with disabilities. For example, the public sector has a number of policies in place that encourage the employment of people with disabilities in a number of different roles, including as teachers, nurses, and social workers.

There are a number of reasons why the public sector has become an important employer of people with disabilities. One reason is that the public sector has a long history of employing people with disabilities. In the 19th century, the public sector employed people with disabilities in a number of different roles, including as clerks, typists, and stenographers. In the 20th century, the public sector employed people with disabilities in a number of different roles, including as teachers, nurses, and social workers.

Another reason why the public sector has become an important employer of people with disabilities is that the public sector has a number of policies in place that encourage the employment of people with disabilities. For example, the public sector has a number of policies in place that encourage the employment of people with disabilities in a number of different roles, including as teachers, nurses, and social workers.

There are a number of reasons why the public sector has become an important employer of people with disabilities. One reason is that the public sector has a long history of employing people with disabilities. In the 19th century, the public sector employed people with disabilities in a number of different roles, including as clerks, typists, and stenographers. In the 20th century, the public sector employed people with disabilities in a number of different roles, including as teachers, nurses, and social workers.

Another reason why the public sector has become an important employer of people with disabilities is that the public sector has a number of policies in place that encourage the employment of people with disabilities. For example, the public sector has a number of policies in place that encourage the employment of people with disabilities in a number of different roles, including as teachers, nurses, and social workers.

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to the quality of the scan and the density of the text. The text appears to be a single paragraph or a series of closely related sentences. The visible fragments of text are as follows:]

[Illegible text block]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to the quality of the scan and the density of the text. The text appears to be a single paragraph or a series of closely related sentences. The visible fragments of text are as follows:]

[Illegible text block]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to the quality of the scan and the density of the text. The text appears to be a single paragraph or a series of closely related sentences. The visible fragments of text are as follows:]

[Illegible text block]

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported to be the most common serotype of *S. flexneri* isolated from children with acute colitis [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1980s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [12]. In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13].

In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13]. In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13].

In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13]. In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13].

In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13]. In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13].

In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13]. In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13].

In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13]. In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13].

In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13]. In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13].

In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13]. In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* from patients with acute colitis in the United Kingdom [13].

the 1990s, the incidence of *S. flexneri* infections has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported as the most common serotype of *Shigella* isolated from children with shigellosis [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1980s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [12]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [13]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [15].

In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [16]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [17]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [18]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [19].

In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [20]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [21]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [22]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [23].

In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [24]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [25]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [26]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [27].

In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [28]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [29]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [30]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [31].

In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [32]. In the 1990s, *S. flexneri* was the most commonly isolated *Shigella* serotype from patients with shigellosis in the United Kingdom [33].

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported to be the most common serotype of *Shigella* isolated from children with shigellosis [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1970s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [12]. In the 1980s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [13].

In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [15].

In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [16]. In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [17].

In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [18]. In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [19].

In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [20]. In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [21].

In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [22]. In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [23].

In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [24]. In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [25].

In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [26]. In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [27].

[The following text is a dense, continuous block of text, likely a scan of a document page. It appears to be a mix of English and possibly some non-English characters, but the overall structure suggests a single paragraph or a series of lines of text. Due to the low resolution and potential noise in the scan, the specific words and punctuation are difficult to discern accurately. The text is oriented vertically on the page.]

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported as the most common serotype in children with acute bacterial dysentery [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1970s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [12]. In the 1980s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [13].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [15].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [16]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [17].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [18]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [19].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [20]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [21].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [22]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [23].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [24]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [25].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [26]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [27].

[The following text is a dense, continuous block of characters and symbols, likely representing a corrupted or heavily redacted document. It contains no legible words or phrases.]

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported as the most common serotype in children with acute bacterial dysentery [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1970s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [12]. In the 1980s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [13].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [14]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [15]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [16].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [17]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [18]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [19].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [20]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [21]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [22].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [23]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [24]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [25].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [26]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [27]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [28].

In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [29]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [30]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [31].

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported as the most common serotype in children with acute bacterial dysentery [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1970s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [12]. In the 1980s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [13]. In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [14].

The aim of this study was to determine the prevalence of *S. flexneri* in children with acute bacterial dysentery in the United Kingdom. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* has increased in the 1990s.

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

[The following text is a dense, continuous block of text, likely a scan of a document page. It is mostly illegible due to extreme blurring and low contrast. The text appears to be a single paragraph or a series of lines of prose, but the specific words and sentences cannot be transcribed accurately.]

